



Towards the Synthesis of Diverse Types of Carrageenan Oligosaccharides

Kinnaert, Christine; Clausen, Mads Hartvig

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Kinnaert, C., & Clausen, M. H. (2017). *Towards the Synthesis of Diverse Types of Carrageenan Oligosaccharides*. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Towards the Synthesis of Diverse Types of Carrageenan Oligosaccharides

Christine Kinnaert¹, Mads Clausen¹

¹Technical University of Denmark. Kgs Lyngby. Denmark.

e-mail: chriki@kemi.dtu.dk

Carrageenans are a family of linear sulfated polysaccharides that are extracted from red seaweeds. They play important roles in the biology of algae and furthermore they have many important industrial applications, such as gelling agent in the food industry. The idealized structures of the 10 known types of carrageenans are presented in Figure 1. The carrageenans are intriguing structures: they have a backbone of D-galactose with alternating β -1,4 and α -1,3 bonds with varying degree of sulfation of the 2, 6, 2' and 4'-positions. Upon treatment with alkali, an 1,3-anhydrogalactose is formed and the α -linked galactose residues undergo a conformational change from 4C_1 to 1C_4 which increases the gelling property of the molecule [2]. We wish to synthesize well-defined oligocarrageenans. Our synthetic strategy aims at producing all ten types of carrageenans from a single precursor carrying different protecting groups that can be cleaved specifically to introduce the sulfate groups at various positions depending on the targeted carrageenan.




Figure 1. Ten types of idealized repeating units of carrageenans [1].

[1] Necas J.; Bartosikova L.; Carrageenan: A review. *Veterinari Medicina* **2013**, 58, 187-205.

[2] Hoffmann R. A.; Gidley M. J.; Cooke D.; Frith W. J.; Effect of isolation procedures on the molecular composition and physical properties of *Eucheuma cottonii* carrageenan. *Food Hydrocolloids* **1995**, 9, 291-289.